

CHILDREN'S ACETAMINOPHEN (TYLENOL)

CHILDREN'S ACETAMINOPHEN (Tylenol)	Infants' Oral Suspension **NEW**	Infants' Concentrated Drops **OLD**	Children's Suspension Liquid	Children's Soft Chews Chewable	Junior Strength Chewable
DOSE**	160 mg/5 mL	80 mg/0.8 mL	160 mg/5 mL	80 mg each	160 mg each
Weight			Teaspoon	Tablet	Tablet
6–11 lbs	1.25 mL	40 mg = (0.4 mL)		--	--
12–17 lbs	2.5 mL	80 mg = (0.8 mL)	½ tsp	--	--
18–23 lbs	3.75 mL	120 mg = (0.8 + 0.4 mL)	¾ tsp	--	--
24–35 lbs	5 mL	160 mg = (0.8 + 0.8 mL)	1 tsp	2 tablets	--
36–47 lbs	--	--	1 ½ tsp	3 tablets	--
48–59 lbs	--	--	2 tsp	4 tablets	2 tablets
60–71 lbs	--	--	2 ½ tsp	5 tablets	2½ tablets
72–95 lbs	--	--	3 tsp	6 tablets	3 tablets
96 lbs & over	--	--	--	--	4 tablets

** All dosages may be repeated every 4 hours, but not more than 5 times daily.

** See box insert for warnings.

CHILDREN'S IBUPROFEN (MOTRIN/ADVIL)

CHILDREN'S IBUPROFEN (Motrin/Advil)	Infants' Concentrated Drops	Children's Suspension	Children's Chewable	Junior Strength Chewable	Junior Strength Caplets
DOSE**	50 mg/1.25 mL	100 mg/5 mL	50 mg	100 mg	100 mg
Weight		Teaspoon	Tablet	Tablet	Caplet
12–17 lbs	50 mg = (1.25 mL)	--	--	--	--
18–23 lbs	75 mg = (1.875 mL)	--	--	--	--
24–35 lbs	--	1 tsp	2 tablets	--	--
36–47 lbs	--	1½ tsp	3 tablets	--	--
48–59 lbs	--	2 tsp	4 tablets	2 tablets	2 caplets
60–71 lbs	--	2½ tsp	5 tablets	2½ tablets	2½ caplets
72–95 lbs	--	3 tsp	6 tablets	3 tablets	3 caplets

** All dosages may be repeated every 6-8 hours, but not more than 4 times daily.

** See box insert for warnings.